

Softly Broken Supersymmetric Gauge Theories through Compactifications

Kazunori Takenaga *

*Department of Physics, Kobe University,
Rokkodai, Nada, Kobe 657, Japan*

Abstract

Effects of boundary conditions of fields for compactified space directions on the supersymmetric gauge theories are discussed. For general and possible boundary conditions the supersymmetry is explicitly broken to yield universal soft supersymmetry breaking terms, and the gauge symmetry of the theory can also be broken through the dynamics of non-integrable phases, depending on number and the representation under the gauge group of matters. The 4-dimensional supersymmetric QCD is studied as a toy model when one of the space coordinates is compactified on S^1 .

*email: takenaga@oct.phys.kobe-u.ac.jp, JSPS Research Fellow

1 Introduction

Our space-time dimensions may be larger than four at fundamental scale such as the Planck scale. Actually, it must be so for consistency of the (super) string theories[1]. In our laboratories, however, we know that our space-time dimensions are four, so that extra space coordinates must be compactified by certain mechanism. Mechanism of the compactifications is still unknown, but physical consequences of compactifications have been studied in various theories since the proposal by Kaluza-Klein[2].

One must specify boundary conditions of fields for compactified directions if space is multiply-connected. We do not know, *a priori*, what boundary conditions should be imposed on the fields for the directions. This is very contrary to the case of the finite temperature field theory, in which the boundary conditions for the euclidean time direction is determined definitely by the quantum statistics of particles.

We shall consider general and possible boundary conditions in supersymmetric gauge theory. One can require that the fields return to their original values up to phases proportional to their charges of global symmetry transformations when the fields travel along the compactified directions[3][4]. The global symmetry transformations must be symmetry of the theory. The lagrangian is automatically single-valued even if the fields have such the boundary conditions.

In a previous paper[5] we studied the effect of the boundary condition associated with the $U(1)_R$ symmetry on the supersymmetry breaking in the supersymmetric QED. The translational invariance for the compactified direction is broken by the boundary condition, so that the variation of action under the supersymmetric transformations does not vanish and remains as surface terms. The supersymmetry is explicitly broken due to the boundary condition. All the effects of the supersymmetry breaking turn out to appear in the lagrangian as the soft supersymmetry breaking terms whose coupling constants are given by an unique parameter and the gauge coupling.

In addition to the boundary condition mentioned above, we can consider the boundary condition associated with the global gauge symmetry in supersymmetric gauge theories. The boundary condition is closely related with the non-integrable phases of the gauge field along the compactified direction, which is dynamical degrees of freedom in multiply-connected space[3][4]. The boundary condition does not break the supersymmetry, but instead, it can break the gauge symmetry of the theory through the dynamics of the non-integrable phases.

In this paper we shall investigate the supersymmetric gauge theory, namely, the supersymmetric QCD (SQCD) when the theory, in which the fields have general and possible boundary conditions for the compactified directions, is compactified. In order to study it as analytically as possible, we shall consider a toy model such that the $SU(N)$ SQCD is compactified on $M^3 \otimes S^1$. In the next section we will define the boundary conditions of

the fields for the S^1 direction. And we will discuss how the boundary condition associated with the $U(1)_R$ symmetry breaks the supersymmetry and how the soft supersymmetry breaking terms appear in the lagrangian. In the section 3 we will give brief summary on the dynamics of the non-integrable phases in multiply-connected space. Then, we will evaluate the effective potential for the non-integrable phases in the SQCD to find how the gauge symmetry is broken. The symmetry breaking depends on number and the representation under the gauge group of matters. The final section is devoted to conclusions and discussions.

2 Soft Supersymmetry Breaking Terms

In this section we shall show how the boundary condition associated with the $U(1)_R$ symmetry breaks the supersymmetry and how the soft supersymmetry breaking terms appear in a toy model, 4-dimensional $SU(N)$ SQCD compactified on $M^3 \otimes S^1$, where M^3 is the 3-dimensional Minkowski space-time, and S^1 is a circle. We use a notation such as $x^{\hat{\mu}} \equiv (x^\mu, x^3) \equiv (x, y)$ and denote the length of the circumference of the S^1 by L .

2.1 Boundary Conditions and Surface Terms

The lagrangian we consider is given by

$$\begin{aligned}\mathcal{L}_{SQCD} &= \frac{1}{2}\text{tr}[W^A W_A + h.c.]_F \\ &= \text{tr}\left[-\frac{1}{2}F_{\hat{\mu}\hat{\nu}}F^{\hat{\mu}\hat{\nu}} - i\lambda\sigma^{\hat{\mu}}D_{\hat{\mu}}\bar{\lambda} + iD_{\hat{\mu}}\lambda\sigma^{\hat{\mu}}\bar{\lambda} + D^2\right].\end{aligned}\quad (1)$$

The subscript F in (1) means F -term. The W_A is the spinorial chiral superfield constructed by the vector superfield V in the Wess-Zumino gauge[6]. The $A(=1, 2)$ stands for a two-component Weyl spinor index. The $W_A(\equiv W_A^a T^a)$ contains the vector boson $A_{\hat{\mu}}$ (gluon), a two-component Weyl fermion λ_A (gaugino) and the auxiliary field D . The auxiliary field is eliminated by the equations of motion for it. The $T^a(a=1, \dots, N^2-1)$ is the generator of $SU(N)$ gauge group. The $F_{\hat{\mu}\hat{\nu}}$ is the field strength for the gluon.

Under the supersymmetric transformations defined by

$$\begin{aligned}\delta_\xi A_{\hat{\mu}} &= \xi\sigma_{\hat{\mu}}\bar{\lambda} - \bar{\xi}\bar{\sigma}_{\hat{\mu}}\lambda, & \delta_\xi \lambda &= \xi D + \sigma^{\hat{\mu}\hat{\nu}}\xi F_{\hat{\mu}\hat{\nu}}, \\ \delta_\xi D &= i\xi\sigma^{\hat{\mu}}D_{\hat{\mu}}\bar{\lambda} + i\bar{\xi}\bar{\sigma}^{\hat{\mu}}D_{\hat{\mu}}\lambda,\end{aligned}$$

the lagrangian varies as $\delta_\xi \mathcal{L}_{SQCD} = \partial_{\hat{\mu}} X^{\hat{\mu}}$, where $X^{\hat{\mu}}$ is calculated as

$$X^{\hat{\mu}} = \left[-\xi\sigma_{\hat{\nu}}\bar{\lambda}^a F^{a\hat{\mu}\hat{\nu}} + \frac{i}{2}\xi\sigma^{\hat{\rho}\hat{\sigma}}\sigma^{\hat{\mu}}\bar{\lambda}^a F_{\hat{\rho}\hat{\sigma}}^a + \frac{i}{2}\xi\sigma^{\hat{\mu}}\bar{\lambda}^a D^a + h.c.\right].$$

The ξ is the supersymmetric transformation parameter of a two-component constant Weyl spinor. The lagrangian (1) is invariant under the $U(1)_R$ transformation defined by

$$W_A(\theta) \rightarrow e^{i\beta} W_A(e^{-i\beta}\theta).$$

The θ is the superspace coordinate. In terms of the component fields, the transformations can be written as

$$\lambda_A \rightarrow e^{i\beta} \lambda_A, \quad A_{\hat{\mu}} \rightarrow A_{\hat{\mu}}, \quad D \rightarrow D,$$

We see that the $U(1)_R$ charges are different between the bosons and the fermions in a supermultiplet W_A .

We define the boundary conditions of the fields for the S^1 -direction as follows;

$$\begin{aligned} A_{\hat{\mu}}(x, y + L) &= U_g A_{\hat{\mu}}(x, y) U_g^\dagger, \\ \lambda(x, y + L) &= e^{i\beta} U_g \lambda(x, y) U_g^\dagger, \\ D(x, y + L) &= U_g D(x, y) U_g^\dagger, \end{aligned} \tag{2}$$

where U_g is a constant $SU(N)$ matrix. The lagrangian (1) is still single-valued even if the fields have such the boundary conditions. If the fields have the boundary condition (2), the surface term,

$$\delta_\xi S_{SQCD} = \left(\int dx \, X^3(x, y) \right) |_{S^1}$$

does not vanish because there is a difference between $X^3(x, y + L)$ and $X^3(x, y)$ due to the non-trivial phase $e^{i\beta}$ in (2). Note that the supersymmetric transformation parameter ξ obeys the periodic boundary condition. The X^3 is the third space-component of the total derivative $X^{\hat{\mu}}$. The translational invariance for the S^1 direction is broken by the boundary condition, so that the supersymmetry is explicitly broken. Note that U_g associated with the global gauge symmetry does not break the translational or supersymmetric invariance.

2.2 Gauged $U(1)_R$ Transformation

For a moment, let us notice the boundary condition associated with the $U(1)_R$ symmetry, that is, the non-trivial phase $e^{i\beta}$ in (2). We shall discuss the importance of U_g on the gauge symmetry breaking in the section 3.

When we expand the fields in the Fourier series for the S^1 direction,

$$\begin{aligned} A_{\hat{\mu}}(x, y) &= \frac{1}{\sqrt{L}} \sum_{n=-\infty}^{+\infty} A_{\hat{\mu}}^{(n)}(x) e^{\frac{2\pi i}{L} n y}, \quad \lambda(x, y) = \frac{1}{\sqrt{L}} \sum_{n=-\infty}^{+\infty} \lambda^{(n)}(x) e^{\frac{2\pi i}{L} (n + \frac{\beta}{2\pi}) y}, \\ D(x, y) &= \frac{1}{\sqrt{L}} \sum_{n=-\infty}^{+\infty} D^{(n)}(x) e^{\frac{2\pi i}{L} n y}, \end{aligned} \tag{3}$$

we observe that $\lambda(x, y)$ given in (3) can be redefined so as to satisfy the periodic boundary condition by gauged $U(1)_R$ transformation whose parameter depends linearly only on the compactified coordinate y [7][8];

$$\lambda(x, y) = U_R(y) \tilde{\lambda}(x, y), \quad U(y)_R \equiv e^{i \frac{\beta}{L} y}, \tag{4}$$

where $\tilde{\lambda}(x, y)$ satisfies $\tilde{\lambda}(x, y + L) = \tilde{\lambda}(x, y)$.

As discussed in the previous paper[5], the supersymmetry breaking terms become manifest in the lagrangian by redefining the fields so as to satisfy the periodic boundary condition. By using (4), $\mathcal{L}_{SQCD}$ can be recast in terms of $(A_{\tilde{\mu}}, \tilde{\lambda})$ as

$$\mathcal{L}_{SQCD} = \tilde{\mathcal{L}}_{SQCD} + \tilde{\mathcal{L}}_{SQCD}^{soft},$$

where $\tilde{\mathcal{L}}_{SQCD}$ has the same form with the original lagrangian except that all the fields satisfy the periodic boundary condition. And $\tilde{\mathcal{L}}_{SQCD}^{soft}$ is obtained as

$$\tilde{\mathcal{L}}_{SQCD}^{soft} = U_R \partial_{\tilde{\mu}} U_R^\dagger \text{tr} [-2i \tilde{\lambda} \sigma^{\tilde{\mu}} \tilde{\lambda}] = -\frac{2\beta}{L} \text{tr} (\bar{\psi}_1 \psi_1 + \bar{\psi}_2 \psi_2),$$

where we have used the Majorana spinors in the 3-dimensions defined by $\tilde{\lambda}_M^T \equiv (\psi_1, i\psi_2)^T$. The $\tilde{\lambda}_M$ is the 4-component Majorana spinors constructed by $\tilde{\lambda}_M^T = (\tilde{\lambda}, \tilde{\tilde{\lambda}})^T$. The supersymmetry breaking terms are the gaugino masses whose coupling constants are given by an unique parameter β . The $\tilde{\mathcal{L}}_{SQCD}^{soft}$ is generated through the derivative in the kinetic term for the gaugino, where the gauged $U(1)_R$ transformation (4) is not respected as symmetry of the theory.

We can define the modified supersymmetry transformations for $(A_{\tilde{\mu}}, \tilde{\lambda})$. The explicit breaking of the supersymmetry due to (2) also becomes manifest by the variation of $\mathcal{L}_{SQCD}$ under the modified supersymmetry transformations. We find

$$\tilde{\delta}_\xi \mathcal{L}_{SQCD} = \partial_{\tilde{\mu}} \tilde{X}^{\tilde{\mu}} + [U_R \partial_{\tilde{\mu}} U_R^\dagger] \tilde{\delta}_\xi \text{tr} [-2i \tilde{\lambda} \sigma^{\tilde{\mu}} \tilde{\lambda}], \quad (5)$$

where $\tilde{\delta}_\xi$ defines the modified supersymmetric transformations. The boundary condition associated with the $U(1)_R$ symmetry breaks the supersymmetry explicitly as shown in the second term in (5). As we expected, the breaking of the supersymmetry is entirely due to the locality of the gauged $U(1)_R$ transformation, *i.e.* $U_R \partial_{\tilde{\mu}} U_R^\dagger$.

2.3 Supersymmetry Breaking Terms from Matters

Let us discuss what types of the supersymmetry breaking terms appear if we add matters. We introduce massive matter superfields $Q_I(\bar{Q}^I)$ which belong to the (anti) fundamental representation under $SU(N)$ gauge group. The $I(= 1, \dots, N_F)$ stands for flavour index. The $Q_I(\bar{Q}^I)$ contains a complex scalar field $\phi_{qI}(\phi_{\bar{q}}^I)$, a two-component Weyl fermion $q_I(\bar{q}^I)$ and the auxiliary field $F_{qI}(F_{\bar{q}}^I)$. In appendix we present the explicit form of the lagrangian for the matters and its variation under the supersymmetry transformations in the 4-dimensions.

The $U(1)_R$ symmetry, which is symmetry of the theory, is defined by

$$Q_I(\theta) \rightarrow e^{i\beta} Q_I(e^{-i\beta} \theta), \quad \bar{Q}^I(\theta) \rightarrow e^{i\beta} \bar{Q}^I(e^{-i\beta} \theta). \quad (6)$$

In terms of the component fields, the transformations can be written as

$$\begin{aligned} \phi_{qI} &\rightarrow e^{i\beta} \phi_{qI}, & q_I &\rightarrow q_I, & F_{qI} &\rightarrow e^{-i\beta} F_{qI}, \\ \phi_{\bar{q}}^I &\rightarrow e^{i\beta} \phi_{\bar{q}}^I, & \bar{q}^I &\rightarrow \bar{q}^I, & F_{\bar{q}}^I &\rightarrow e^{-i\beta} F_{\bar{q}}^I. \end{aligned} \quad (7)$$

We see that the $U(1)_R$ charges are different between the bosons and the fermions in each supermultiplet $Q_I, \bar{Q}^I$.

The boundary conditions we take are

$$\begin{aligned}\phi_{qI}(x, y+L) &= e^{i\beta} U_g \phi_{qI}(x, y), & \phi_q^I(x, y+L) &= e^{i\beta} \bar{U}_g \phi_q^I(x, y), \\ q_I(x, y+L) &= U_g q_I(x, y), & \bar{q}^I(x, y+L) &= \bar{U}_g \bar{q}^I(x, y), \\ F_{qI}(x, y+L) &= e^{-i\beta} U_g F_{qI}(x, y), & F_q^I(x, y+L) &= e^{-i\beta} \bar{U}_g F_q^I(x, y),\end{aligned}\tag{8}$$

where $U_g, \bar{U}_g$ are constant $SU(N), SU(\bar{N})$ matrices, respectively. Their importance on the gauge symmetry breaking are discussed in the next section. Under the supersymmetry transformations, the lagrangian varies as $\delta_\xi \mathcal{L}_{matters} = \partial_{\hat{\mu}} X_{matters}^{\hat{\mu}}$. The third space component of the total derivative $X_{matters}^3$ do not return to their original values after the translation along the S^1 -direction due to the non-trivial phase in (8). Therefore, the supersymmetry is explicitly broken.

When we expand the fields in the Fourier series for the S^1 -direction, we find

$$\begin{aligned}\phi_{qI}(x, y) &= U_R(y) \tilde{\phi}_{qI}(x, y), & \phi_q^I(x, y) &= U_R(y) \tilde{\phi}_q^I(x, y), \\ F_{qI}(x, y) &= U_R^\dagger(y) \tilde{F}_{qI}(x, y), & F_q^I(x, y) &= U_R^\dagger(y) \tilde{F}_q^I(x, y).\end{aligned}\tag{9}$$

As before, the fields can be redefined so as to satisfy the periodic boundary condition by the gauged $U(1)_R$ transformation $U_R(y) = e^{i\frac{\beta}{L}y}$. The $\tilde{\phi}_{qI}, \tilde{\phi}_q^I, \tilde{F}_{qI}$ and $\tilde{F}_q^I$ satisfy the periodic boundary condition.

The supersymmetry breaking terms manifestly appear by rewriting the lagrangian in terms of the fields with the periodic boundary condition. By using (9), we obtain

$$\mathcal{L}_{SQCD} + \mathcal{L}_{matters} = \tilde{\mathcal{L}}_{SQCD} + \tilde{\mathcal{L}}_{matters} + \tilde{\mathcal{L}}^{soft},$$

where $\tilde{\mathcal{L}}_{SQCD}$ and $\tilde{\mathcal{L}}_{matters}$ are the same form with the original lagrangian except that all the fields satisfy the periodic boundary condition. The $\tilde{\mathcal{L}}^{soft}$ is generated through the derivatives in the kinetic terms for the gaugino and the squark, where the gauged $U(1)_R$ transformation by $U_R(y)$ is not respected as symmetry of the theory. The supersymmetry breaking terms are never generated from the superpotential $W(Q, \bar{Q})$ because there are no derivatives in it. The $\tilde{\mathcal{L}}^{soft}$ is obtained as

$$\begin{aligned}\tilde{\mathcal{L}}^{soft} &= U_R(y) \partial_{\hat{\mu}} U_R^\dagger(y) \left[\text{tr}(-2i\tilde{\lambda} \sigma^{\hat{\mu}} \tilde{\lambda}) + \sum_I \left(\tilde{\phi}_{qI}^\dagger (D^{\hat{\mu}} \tilde{\phi}_{qI}) - (D^{\hat{\mu}} \tilde{\phi}_{qI})^\dagger \tilde{\phi}_{qI} \right) \right. \\ &\quad + \sum_I \left(\tilde{\phi}_q^{I\dagger} (D^{\hat{\mu}} \tilde{\phi}_q^I) - (D^{\hat{\mu}} \tilde{\phi}_q^I)^\dagger \tilde{\phi}_q^I \right) \left. \right] + |\partial_{\hat{\mu}} U_R(y)|^2 \sum_I \left(|\tilde{\phi}_{qI}|^2 + |\tilde{\phi}_q^I|^2 \right) \\ &\longrightarrow - \frac{2\beta}{L} \text{tr}(\bar{\psi}_1 \psi_1 + \bar{\psi}_2 \psi_2) + \left(\frac{\beta}{L} \right)^2 \sum_I \left(|\tilde{\phi}_{qI}|^2 + |\tilde{\phi}_q^I|^2 \right) \\ &\quad + \frac{2\beta g}{L} \sum_I \left(\tilde{\phi}_{qI}^\dagger T^a \tilde{\phi}_{qI} - \tilde{\phi}_q^{I\dagger} T^a \tilde{\phi}_q^I \right) \Phi^a.\end{aligned}\tag{10}$$

The g is the gauge coupling constant. The $\rightarrow$ in (10) means that the dimensional reduction from $D = 4$ to $D = 3$ is carried out, ignoring the Kaluza-Klein modes ($n \neq 0$) in the Fourier series. We have denoted the gauge field for the S^1 -direction A_3^a as Φ^a . The Φ^a is a real scalar field which belongs to the adjoint representation under the gauge group. We realize again that the supersymmetry breaking is entirely due to the locality of the gauged $U(1)_R$ transformation, *i.e.* $U_R \partial_{\hat{\mu}} U_R^\dagger$ and $|\partial_{\hat{\mu}} U_R|^2$.

We find that the supersymmetry breaking terms are the scalar mass and the trilinear scalar terms whose couplings depend only on a unique parameter β and the gauge coupling g . As the remarkable consequence, the supersymmetry breaking terms generated in this mechanism are common to all flavours and are soft breaking. This is because the derivative ∂_μ are common to all flavours and has mass dimension one, so that the couplings generated through the derivative are always universal and dimensional couplings. The universality may be needed to avoid the FCNC.

We also mention the supertrace of the squared mass matrix $\text{Str} \mathcal{M}^2 = \sum_J (-)^{2J} (2J + 1) \mathcal{M}_J^2 = 0$, where J stands for the spin of the particles. In our case the masses for the gaugino and the quarks are β/L and m , respectively. On the other hand, the mass for the squarks is $m^2 - (\beta/L)^2$. The gluon A_μ^a and the scalar Φ^a are massless. It is evaluated as[9]

$$\text{Str} \mathcal{M}^2 = 2(m^2 - (\beta/L)^2) - 2(m^2 + (\beta/L)^2) = -4(\beta/L)^2 \neq 0. \quad (11)$$

This result is expected because the supersymmetry is broken explicitly in our case. This may be desirable when we try to build models with the soft supersymmetry breaking terms based on our mechanism.

One may think that the boundary condition associated with the global flavour symmetry is possible. The boundary condition, however, does not break the supersymmetry because their charges of global flavour transformations are the same between the bosons and the fermions in a supermultiplet. In this case we would obtain supersymmetric invariant soft terms generated by the same manner discussed in this section.

3 Non-integrable Phase and Its Dynamics in SQCD

We have discussed that the boundary condition associated with the $U(1)_R$ symmetry breaks the supersymmetry. In this section we shall discuss the role of the boundary condition associated with the global gauge symmetry, which have been ignored in the previous section. Readers familiar with this topics can skip argument below and go to the subsection 3.1 directly. Details discussions are given in[4].

For general discussions we shall consider $SU(N)$ SQCD on $M^{D-n} \otimes T^n$ in this section, where M^{D-n} and T^n are $(D-n)$ -dimensional Minkowski space-time and n -torus, respectively. The $x^\mu (\mu = 0, 1, \dots, D-(n+1))$ are the coordinates of M^{D-n} , and $y^a (a = 1, \dots, n)$

are those of T^n . We use a notation $x^{\hat{\mu}}$ (or x) which stands for $x^{\hat{\mu}} \equiv x \equiv (x^\mu, y^a)$. We define the boundary conditions as follows;

$$\begin{aligned} A_{\hat{\mu}}(x^\mu, \dots y^a + L^a \dots) &= U_g^a A_{\hat{\mu}}(x^\mu, \dots y^a \dots) U_g^{a\dagger}, \\ \lambda(x^\mu, \dots y^a + L^a \dots) &= e^{i\beta} U_g^a \lambda(x^\mu, \dots y^a \dots) U_g^{a\dagger}. \end{aligned} \quad (12)$$

The $L^a (a = 1, \dots, n)$ is the length of the circumference of each circle, and a constant matrix $U_g^a (a = 1 \dots n) \in SU(N)$. The lagrangian is still single-valued with the fields having the boundary conditions. We symbolically denote the boundary condition as $U_g^a (a = 1, \dots, n)$. In order to fix a theory with the boundary conditions (12) definitely, a set of $U_g^a (a = 1, \dots, n)$ has given for any configurations of the fields $A_{\hat{\mu}}, \lambda$.

There exists a class of gauge transformation which does not change the boundary condition U_g^a . We denote the class of the gauge transformation by $\Omega(x)$. Under the gauge transformation $\Omega(x)$, the fields transform as

$$\begin{aligned} A_{\hat{\mu}}(x) \rightarrow A'_{\hat{\mu}}(x) &= \Omega(x) A_{\hat{\mu}}(x) \Omega^\dagger(x) + \frac{i}{g} \Omega(x) \partial_{\hat{\mu}} \Omega^\dagger(x) \\ \lambda(x) \rightarrow \lambda'(x) &= \Omega(x) \lambda(x) \Omega^\dagger(x). \end{aligned}$$

The boundary conditions for the transformed fields $A'_{\hat{\mu}}, \lambda'$ are the same with those for $A_{\hat{\mu}}, \lambda$. Hence, we have

$$U_g^a = \Omega(x^\mu, y^a + L^a) U_g^a \Omega^\dagger(x^\mu, y^a). \quad (13)$$

We can consider another class of the gauge transformation by which the boundary condition is changed. Let us denote such gauge transformation by $\mathcal{T}(x)$. Under the transformation by $\mathcal{T}(x)$, the fields are redefined in the form of the gauge transformation by

$$\begin{aligned} A'_{\hat{\mu}}(x) &= \mathcal{T}(x) A_{\hat{\mu}}(x) \mathcal{T}^\dagger(x) + \frac{i}{g} \mathcal{T}(x) \partial_{\hat{\mu}} \mathcal{T}^\dagger(x), \\ \lambda'(x) &= \mathcal{T}(x) \lambda(x) \mathcal{T}^\dagger(x). \end{aligned}$$

In order for the redefined fields $A'_{\hat{\mu}}, \lambda'$ to satisfy (12) with new boundary condition $U_g^{a'}$, we have

$$U_g^{a'} = \mathcal{T}(x^\mu, y^a + L^a) U_g^a \mathcal{T}^\dagger(x^\mu, y^a). \quad (14)$$

The gauge transformations $\mathcal{T}(x)$ (and $\Omega(x)$) must satisfy

$$\partial_{\hat{\mu}} [\mathcal{T}(x^\mu, y^a + L^a) U_g^a \mathcal{T}^\dagger(x^\mu, y^a)] = 0$$

to maintain the x -independence of (new) boundary condition.

Let us assume $\langle F_{\hat{\mu}\hat{\nu}} \rangle = 0$ in the vacuum, and it follows that

$$\langle A_{\hat{\mu}} \rangle = \frac{i}{g} V^\dagger(x) \partial_{\hat{\mu}} V(x). \quad (15)$$

It is important to note that this pure gauge configuration, in general, is physically distinct from $\langle A_{\hat{\mu}} \rangle = 0$ in multiply-connected space because the transformation with $\mathcal{T}(x)$ change the boundary condition as seen from (14). The $V(x)$ is determined by quantum effects as a function of U_g^a up to a global gauge transformation. If we perform the gauge transformation with $\mathcal{T}(x) = V(x)$, then we have $\langle A'_{\hat{\mu}} \rangle = 0$ with new boundary condition such as

$$U_g^{a:inv} \equiv U_g^{a'} = V(x^\mu, y^a + L^a) U_g^a V^\dagger(x^\mu, y^a). \quad (16)$$

Since $U_g^{a:inv}$ is the boundary condition for $\langle A'_{\hat{\mu}} \rangle = 0$, the symmetry of the theory is generated by the generators of the gauge group which commute with $U_g^{a:inv}$. The gauge transformation which satisfy

$$\Omega(x^\mu, y^a + L^a) = U_g^{a:inv} \Omega(x^\mu, y^a) U_g^{a:inv\dagger}(x^\mu, y^a) \quad (17)$$

is large gauge transformation under which the non-integrable phases are invariant. We will find that this invariance is reflected in the effective potential for the non-integrable phases.

Let us show that $U_g^{a:inv}$ is closely related with the path-ordered integral along a loop C of the compactified coordinate;

$$W_c^a(C) \equiv \mathcal{P} \exp[-ig \int_C dy^a A_a] U_g, \quad (a = 1, \dots, n).$$

We can define n numbers of path-ordered integrals for each gauge field $A_a (a = 1, \dots, n)$ along each compactified coordinate $y^a (a = 1, \dots, n)$. For the pure gauge configuration (15), it is evaluated as

$$W_c^a(C) = V^\dagger(x^\mu, y^a) V(x^\mu, y^a + L^a) U_g^a. \quad (18)$$

Hence, from (16), $W_c^a(C)$ is related to $U_g^{a:inv}$ as

$$W_c^a(C) = V^\dagger(x^\mu, y^a) U_g^{a:inv} V(x^\mu, y^a).$$

The eigenvalues of $U_g^{a:inv}$ or $W_c^a(C)$ are called non-integrable phases, which are important quantities in discussing the gauge symmetry breaking of the theory.

One of the boundary conditions $U_g^a (a = 1, \dots, n)$ for the gauge field A_a can be diagonalized by utilizing the degrees of freedom of global gauge transformation. Here we assume that all of them have a diagonal form as a special case;

$$U_g^a = \begin{pmatrix} e^{i\lambda_{a,1}} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & e^{i\lambda_{a,N}} \end{pmatrix}, \quad \sum_{i=1}^N \lambda_{a,i} = 0, \quad (a = 1, \dots, n), \quad (19)$$

where $i(= 1, \dots, N)$ stands for the indices of $SU(N)$ gauge group. Let us perform the gauge transformation $\mathcal{T}(y)$ given by

$$\mathcal{T}(y) = \begin{pmatrix} e^{i\varphi_1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & e^{i\varphi_N} \end{pmatrix}, \quad \text{with} \quad \varphi_i \equiv \sum_{b=1}^n \frac{h_{b,i}}{L^b} y^b, \quad \sum_{i=1}^N h_{b,i} = 0. \quad (20)$$

Under the gauge transformation with (20), the boundary condition (19) changes to

$$\begin{aligned} U_g^{a'} &= \mathcal{T}(y^a + L^a) U_g^a \mathcal{T}^\dagger(y^a) \\ &= \begin{pmatrix} e^{i(\lambda_{a,1} + h_{a,1})} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & e^{i(\lambda_{a,N} + h_{a,N})} \end{pmatrix}. \end{aligned} \quad (21)$$

We can redefine the fields so as to satisfy the periodic boundary condition by choosing $\lambda_{a,i} = -h_{a,i}$. New gauge field with $U_g^{a'} = \mathbf{1}_{N \times N}$ is redefined as

$$A'_{\hat{\mu}} = \mathcal{T} A_{\hat{\mu}} \mathcal{T}^\dagger + \frac{i}{g} \mathcal{T} \partial_{\hat{\mu}} \mathcal{T}^\dagger = \mathcal{T} A_{\hat{\mu}} \mathcal{T}^\dagger + \delta_{\hat{\mu},a} \frac{1}{gL^a} \begin{pmatrix} h_{a,1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & h_{a,N} \end{pmatrix}. \quad (22)$$

We see that the effect of the boundary condition shifts the gauge field by the constant, the second term in (22). New gauge field $A'_a(a = 1, \dots, n)$ satisfies the periodic boundary condition $U_g^{a'} = \mathbf{1}_{N \times N}$, so that we expect $\langle A'_a \rangle$ to be constant. Hence, we parameterize it as follows;

$$\begin{aligned} \langle A'_\mu \rangle &= 0, \quad (\mu = 0, 1, \dots, D - (n + 1)) \\ \langle A'_a \rangle &= \frac{1}{gL^a} \begin{pmatrix} \theta_{a,1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & \theta_{a,N} \end{pmatrix}, \quad \sum_{i=1}^N \theta_{a,i} = 0. \end{aligned} \quad (23)$$

The constant background gauge field (23) can be written in the pure gauge form;

$$\langle A'_a \rangle = \frac{i}{g} V'^\dagger(y) \partial_a V'(y), \quad (24)$$

where $V'(y)$ is given by

$$V'(y) = \begin{pmatrix} e^{-i \sum_{b=1}^n \frac{\theta_{b,1}}{L^b} y^b} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & e^{-i \sum_{b=1}^n \frac{\theta_{b,N}}{L^b} y^b} \end{pmatrix} \in SU(N).$$

Under the gauge transformation with $V'(y)$, new boundary condition for $\langle A_a'' \rangle = 0$ is obtained as

$$U_g^{a''} = V'(y^a + L^a)U'_g V'^{\dagger}(y^a) = \begin{pmatrix} e^{-i\theta_{a,1}} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & e^{-i\theta_{a,N}} \end{pmatrix} \equiv U_g^{a:inv}. \quad (25)$$

On the other hand, the path-ordered integral for (24) is evaluated as

$$W_c^a(C) = V'^{\dagger}(x^\mu, y^a) V'(x^\mu, y^a + L^a) U_g^{a'} = V'^{\dagger}(x^\mu, y^a) U_g^{a:inv} V'(x^\mu, y^a),$$

where we have used (25). The $\theta_{a,i}$ ($a = 1, \dots, n$; $i = 1, \dots, N$) is the eigenvalue of W_c^a or $U_g^{a:inv}$, that is, non-integrable phase.

One can return to the original gauge field $\langle A_a \rangle$ from (24) by the transformation with $\mathcal{T}^{-1}(y)$. We would also obtain the same $U_g^{a:inv}$ with (25) in this gauge.

3.1 Effective Potential for Non-integrable Phases in SQCD

Now, let us discuss the dynamics of the non-integrable phases, which are dynamical degrees of freedom and can not be gauged away in a multiply-connected space. We shall compute the effective potential for the non-integrable phases in the SQCD. It is worth noting that at the tree-level the effective potential does not depend on the non-integrable phases, and the vacuum has the continuous degeneracy. The quantum effects, however, lift the degeneracy. We shall use the perturbation theory in one-loop approximation based on the background field method in the Feynman gauge. The effective potential for the non-integrable phases have been evaluated in non-supersymmetric gauge theories [3][4][10][11][12].

We understand that the effects of the boundary condition associated with the global gauge symmetry are interpreted as the constant background gauge field with the periodic boundary condition[13]

$$\langle A_a \rangle = \frac{1}{gL^a} \begin{pmatrix} \theta_{a,1} & & & \\ & \ddots & & \\ & & \ddots & \\ & & & \theta_{a,N} \end{pmatrix}, (a = 1, \dots, n) \quad \text{with} \quad \sum_{i=1}^N \theta_{a,i} = 0. \quad (26)$$

The $\theta_{a,i}$ is the eigenvalue of the non-integrable phase which we shall determine dynamically. The field strength for (26) vanishes, $\langle F_{\hat{\mu}\hat{\nu}} \rangle = 0$. According to the prescription of the background field method, we obtain the effective potential for the non-integrable phases in the D -dimensional SQCD as follows;

$$\begin{aligned} V_{eff}^{SQCD} &= V_{g+gh} + V_{gaugino} \\ &= -\frac{i}{2}(D-2) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}] + \frac{i}{2}(r2^{\lfloor \frac{D}{2} \rfloor}) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}]. \end{aligned} \quad (27)$$

The $V_{g+gh}, V_{gaugino}$ stand for the contributions from gauge-ghost, gaugino, respectively. The $D - 2$ in (27) is the number of on-shell bosonic degrees of freedom, and $r2^{[D/2]}$ is the one of fermionic degrees of freedom, where the factor $r = 1/2$ if the gaugino λ is a Majorana or Weyl, and $r = 1/4$ if λ is a Majorana-Weyl.

We can expand the fields in the Fourier series for T^n -direction as

$$\begin{aligned} [A_{\hat{\mu}}(x)]_{ij} &= \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \sum_{n_1 \dots n_n} [A_{\hat{\mu}}^{(n_1 \dots n_n)}(x^\mu)]_{ij} \exp\left(\sum_{a=1}^n \frac{2\pi i}{L^a} n_a y^a\right), \\ [\lambda(x)]_{ij} &= \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \sum_{n_1 \dots n_n} [\lambda^{(n_1 \dots n_n)}(x^\mu)]_{ij} \exp\left(\sum_{a=1}^n \frac{2\pi i}{L^a} \left(n_a + \frac{\beta}{2\pi}\right) y^a\right), \end{aligned} \quad (28)$$

where i, j are indices of $SU(N)$ gauge group. The effect of the boundary condition associated with the $U(1)_R$ symmetry manifestly appear in the momentum of the compactified direction as $p_a = \frac{2\pi}{L^a} \left(n_a + \frac{\beta}{2\pi}\right)$ ($a = 1, \dots, n$) for the gaugino field. The covariant derivative $D_{\hat{\mu}}$ in (27) is the one with the constant background gauge field (26), which is evaluated as

$$\begin{aligned} [D_{\hat{\mu}} D^{\hat{\mu}}]_{ij} &= D_\mu D^\mu + D_a D^a, \quad (\mu = 0, 1, \dots, D - (n + 1); a = 1, \dots, n) \\ &= \partial_\mu^2 - D_a^2 \\ &= \partial_\mu^2 - \sum_{a=1}^n \left[\partial_a - \frac{i}{L^a} (\theta_{a,i} - \theta_{a,j})\right]^2. \end{aligned}$$

Then, (27) becomes

$$\begin{aligned} V_{eff}^{SQCD} &= \frac{1}{2} (D - 2) \sum_{n_1 \dots n_n} \sum_{i,j=1}^N \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \\ &\times \int \frac{d^{D-n} p}{(2\pi)^{D-n}} \ln \left[p^2 + \sum_{a=1}^n \left(\frac{2\pi}{L^a}\right)^2 \left(n_a - \frac{\theta_{a,i} - \theta_{a,j}}{2\pi}\right)^2 \right]_{ij} \\ &- \frac{1}{2} r 2^{[D/2]} \sum_{n_1 \dots n_n} \sum_{i,j=1}^N \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \\ &\times \int \frac{d^{D-n} p}{(2\pi)^{D-n}} \ln \left[p^2 + \sum_{a=1}^n \left(\frac{2\pi}{L^a}\right)^2 \left(n_a - \frac{\theta_{a,i} - \theta_{a,j} - \beta}{2\pi}\right)^2 \right]_{ij}. \end{aligned} \quad (29)$$

The off-diagonal ($i \neq j$) components contribute to the effective potential.

It is clear that if $\beta = 0$, the effective potential vanishes for any values of θ 's in the space-time dimensions $D = 3, 4, 6, 10$ because of the equal number of physical degrees of freedom between the bosons and the fermions in the supersymmetric Yang-Mills theory[16];

$$(D - 2) - r 2^{[D/2]} = 0 \quad \text{for} \quad \begin{cases} D = 3, & r = \frac{1}{2}, \\ D = 4, & r = \frac{1}{2}, \\ D = 6, & r = \frac{1}{2}, \\ D = 10, & r = \frac{1}{4}. \end{cases}$$

The $\beta = 0$ restores the supersymmetry to yield the vanishing effective potential for any values of θ 's due to the non-renormalization theorem in the supersymmetric theories. The eigenvalues of the non-integrable phases can not be determined dynamically in this case. The boundary condition associated with the $U(1)_R$ symmetry explicitly breaks the supersymmetry to yield non-vanishing effective potential. We can, in principle, determine the non-integrable phases dynamically as the minimum of the effective potential (29).

3.1.1 Pure SQCD

In order to demonstrate dynamical determination of the non-integrable phases as analytically as possible, we consider a toy model, 4-dimensional SQCD compactified on $M^3 \otimes S^1$. In this case the boundary conditions we take are reduced from (12) to

$$A_{\hat{\mu}}(x^\mu, y + L) = U_g A_{\hat{\mu}}(x^\mu, y) U_g^\dagger, \quad \lambda(x^\mu, y + L) = e^{i\beta} U_g \lambda(x^\mu, y) U_g^\dagger. \quad (30)$$

The effective potential is evaluated as

$$V_{eff}^{SQCD} = -(4 - 2) \frac{\Gamma(2)}{\pi^2 L^4} \sum_{i,j=1}^N \sum_{n=1}^{\infty} \frac{1}{n^4} \cos(n(\theta_i - \theta_j)) + \frac{1}{2} 2^2 \frac{\Gamma(2)}{\pi^2 L^4} \sum_{n=1}^{\infty} \frac{1}{n^4} \cos(n(\theta_i - \theta_j - \beta)). \quad (31)$$

We keep only θ -dependent terms after the momentum integrations in (29). Moreover, if we assume that the gauge group is $SU(2)$, we have only one order parameter $\theta_1 = -\theta_2 \equiv \theta$. The summation with respect to the Fourier mode n can be done by the formula;

$$F(t) \equiv \sum_{n=1}^{\infty} \frac{1}{n^4} \cos nt = -\frac{1}{48} t^2 (t - 2\pi)^2 + \frac{\pi^4}{90} \quad \text{for } 0 \leq t \leq 2\pi. \quad (32)$$

Using this formula, we obtain

$$\tilde{V}_{eff}^{SQCD} \equiv \frac{V_{eff}^{SQCD}}{2/\pi^2 L^4} = -2F(t) + [F(t - \beta) + F(t + \beta)] \equiv -2F(t) + G(t, \beta), \quad (33)$$

where we have defined $t \equiv 2\theta$ and $G(t, \beta) \equiv F(t - \beta) + F(t + \beta)$. The $F(t)$ stands for the gauge and ghost contributions, and $G(t, \beta)$ stands for the gaugino contribution.

The $F(t)$ and $G(t, \beta)$ have properties such that $F(t) = F(-t) = F(t + 2\pi m)$ and $G(t, \beta) = G(-t, \beta) = G(t, -\beta) = G(t + 2\pi m, \beta) = G(t, \beta + 2\pi m)$, respectively. The m is integer. The effective potential has the periodicity $V_{eff}^{SQCD}(t, \beta) = V_{eff}^{SQCD}(t + 2\pi m, \beta)$. The periodicity of the effective potential is traced back to the large gauge transformation generated by $\Omega(x)$ in (17). There is also the periodicity such as $V_{eff}^{SQCD}(t, \beta) = V_{eff}^{SQCD}(t, \beta + 2\pi m)$. This periodicity follows from the redefinition of the gaugino field $\lambda \rightarrow e^{2\pi m y/L} \lambda$.

By straightforward calculations, we find

$$\tilde{V}_{eff}^{SQCD} = -\frac{\beta^2}{12} \left[3t^2 - 6\pi t + 2\pi^2 + \frac{1}{2} \beta^2 \right] \geq \tilde{V}_{eff}(t = 0 \pmod{2\pi}, \beta).$$

The eigenvalues of the non-integrable phases are determined dynamically at $\theta = 0 \pmod{\pi}$ as the minimum of the effective potential. In this case the gauge symmetry is not broken because $U_g^{inv} = \mathbf{1}_{2 \times 2}$ commutes with all the generators of $SU(2)$ gauge group. Note that the parameter β is not the order parameter of the effective potential.

3.1.2 Matters in the Fundamental Representation

If we add massive matters $Q(\phi_q, q)$ and $\bar{Q}(\phi_{\bar{q}}, \bar{q})$, then, additional contributions to the effective potential arise from quarks $q, \bar{q}$ and squarks $\phi_q, \phi_{\bar{q}}$. The $U(1)_R$ symmetry are defined by (6), from which we see that the squarks have the $U(1)_R$ charges, but the quarks do not have them. The boundary conditions of these fields are defined by (8).

The effective potential for the non-integrable phases arising from the (s)quarks is[14]

$$\begin{aligned} V_{eff}^{fd} &= V_{squarks}^{fd} + V_{quarks}^{fd} \\ &= -\frac{i}{2}(2N_F \times 2) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}] + \frac{i}{2} 2N_F \times (r 2^{\lfloor \frac{D}{2} \rfloor}) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}], \end{aligned} \quad (34)$$

where N_F is number of the flavour. The superfields $Q, (\bar{Q})$ belong to the (anti) fundamental representation under $SU(2)$ gauge group. The covariant derivative $D_{\hat{\mu}}$ in (34) is evaluated for these representations as

$$\begin{aligned} [D_{\hat{\mu}} D^{\hat{\mu}}]_{ij} &= D_{\mu} D^{\mu} + D_a D^a, \quad (i = 0, 1, \dots, D - (n + 1); a = 1, \dots, n) \\ &= \partial_{\mu}^2 - D_a^2 \\ &= \left[\partial_{\mu}^2 - \sum_{a=1}^n [\partial_a - \frac{i}{L^a} \theta_{a,i}]^2 \right] \delta_{ij}. \end{aligned}$$

Hence, we obtain

$$\begin{aligned} V_{eff}^{fd} &= \frac{1}{2} 2N_F \times 2 \sum_{n_1 \dots n_n} \sum_{i=1}^N \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \\ &\times \int \frac{d^{D-n} p}{(2\pi)^{D-n}} \ln \left[p^2 + \sum_{a=1}^n \left(\frac{2\pi}{L^a} \right)^2 \left(n_a - \frac{\theta_{a,i} - \beta}{2\pi} \right)^2 \right] \\ &- \frac{1}{2} 2N_F \times r 2^{\lfloor D/2 \rfloor} \sum_{n_1 \dots n_n} \sum_{i,j=1}^N \frac{1}{\sqrt{\prod_{a=1}^n L^a}} \\ &\times \int \frac{d^{D-n} p}{(2\pi)^{D-n}} \ln \left[p^2 + \sum_{a=1}^n \left(\frac{2\pi}{L^a} \right)^2 \left(n_a - \frac{\theta_{a,i}}{2\pi} \right)^2 \right]. \end{aligned}$$

In the 4-dimensions, this yields, keeping only θ -dependent terms after the momentum integrations,

$$V_{eff}^{fd} = -2N_F \times 2 \frac{\Gamma(2)}{\pi^2 L^4} \sum_{n=1}^{\infty} \sum_{i=1}^N \frac{1}{n^4} \cos(n(\theta_i - \beta)) + 2N_F \times \frac{1}{2} 2^2 \frac{\Gamma(2)}{\pi^2 L^4} \sum_{n=1}^{\infty} \sum_{i=1}^N \frac{1}{n^4} \cos(n\theta_i),$$

Then, for $SU(2)$ case, we obtain

$$\begin{aligned}\tilde{V}_{eff}^{fd} &\equiv \frac{V_{eff}^{fd}}{2/\pi^2 L^4} = N_F \left[-2 \times \left[F(t/2 - \beta) + F(t/2 + \beta) \right] + 2 \times 2 \times F(t/2) \right] \\ &\equiv N_F \left[-2G(t/2, \beta) + 2 \times 2 \times F(t/2) \right] \quad \text{for } 0 \leq t/2 \leq 2\pi.\end{aligned}$$

The $G(t/2, \beta) \equiv F(t/2 - \beta) + F(t/2 + \beta)$, $F(t/2)$ stands for the contributions from the squarks, quarks, respectively. As before, if $\beta = 0$, the supersymmetry is restored, and $\tilde{V}_{eff}^{fd}$ vanishes due to the non-renormalization theorem.

Adding these flavour contributions to the effective potential V_{eff}^{SQCD} , we obtain

$$\begin{aligned}\tilde{V}_{eff}^{SQCD+fd} &= \left[-2F(t) + G(t, \beta) \right] + N_F \left[-2G(t/2, \beta) + 2 \times 2 \times F(t/2) \right] \\ &= \frac{\beta^2}{12} \left[-3t^2 + 6\pi t - 2\pi^2 - \frac{1}{2}\beta^2 + N_F \left(\frac{3}{2}t^2 - 6\pi t + 4\pi^2 + \beta^2 \right) \right].\end{aligned}$$

The $-2F(t) + G(t, \beta)$, which comes from the gluon and the gaugino, has minima at $\theta = 0 \pmod{\pi}$. On the other hand, $-2G(t/2, \beta) + 2 \times 2 \times F(t/2)$, which comes from the (s)quark, has minima at $\theta = \pi \pmod{2\pi}$. Hence, $\tilde{V}_{eff}^{SQCD+fd}$ has minima at $\theta = \pi \pmod{2\pi}$ independent of N_F , so that $U_g^{inv} = -\mathbf{1}_{2 \times 2}$. The $SU(2)$ gauge symmetry is not broken for any numbers of the flavour in the fundamental representation.

3.1.3 Matters in the Adjoint Representation

Next, let us add massive matters in the adjoint representation under $SU(2)$ gauge group instead of those in the fundamental representation. We denote the superfield as Q^{adj} which contain adjoint-quark q_I^{adj} and adjoint-squark ϕ_{qI}^{adj} . The $U(1)_R$ symmetry is defined as the same way as the matters in the fundamental representation. We take the boundary conditions of the fields as

$$q_I^{adj}(x, y + L) = U_g q_I^{adj}(x, y) U_g^\dagger, \quad \phi_{qI}^{adj}(x, y + L) = e^{i\beta} U_g \phi_{qI}^{adj}(x, y) U_g^\dagger. \quad (35)$$

The effective potential for the non-integrable phases arising from the adjoint-(s)quark is

$$\begin{aligned}V_{eff}^{adj} &= V_{squarks}^{adj} + V_{quarks}^{adj} \\ &= -\frac{i}{2} (N_F^{adj} \times 2) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}] + \frac{i}{2} N_F^{adj} \times (r 2^{\lfloor \frac{D}{2} \rfloor}) \int_k \text{Tr} \ln [D_{\hat{\mu}} D^{\hat{\mu}}], \quad (36)\end{aligned}$$

The computation goes the same as before. We obtain

$$\begin{aligned}\tilde{V}_{eff}^{adj} &\equiv \frac{V_{eff}^{adj}}{2/\pi^2 L^4} = N_F^{adj} \left[-\left[F(t - \beta) + F(t + \beta) \right] + 2F(t) \right] \\ &\equiv N_F^{adj} \left[-G(t, \beta) + 2F(t) \right] \quad \text{for } 0 \leq t \leq 2\pi.\end{aligned}$$

The $-G(t, \beta)$, $2F(t)$ stand for the squark, quark contributions, respectively. This result also corresponds to massless limit of the matters as before. The $\beta = 0$ restores the

supersymmetry to yield vanishing effective potential again. The total effective potential becomes

$$\begin{aligned}\tilde{V}_{eff}^{SQCD+adj} &= [-2F(t) + G(t, \beta)] + N_F^{adj} [-G(t, \beta) + 2 \times F(t)] \\ &= \frac{\beta^2}{12} \left[-3t^2 + 6\pi t - 2\pi^2 - \frac{1}{2}\beta^2 + N_F^{adj} \left(3t^2 - 6\pi t + 2\pi^2 + \frac{1}{2}\beta^2 \right) \right].\end{aligned}$$

For $N_F^{adj} \neq 1$, we find the minimum of the potential is located at $\theta = \pi/2$ independent of N_F^{adj} . This means $U_g^{inv} = -i\sigma^3$, so that the gauge symmetry is broken to $U(1)$. Even if we take $N_F^{adj} \rightarrow \infty$, the gauge symmetry is never restored to $SU(2)$. This is very contrary to the case for the matters in the fundamental representation.

For $N_F^{adj} = 1$, the effective potential vanishes for any values of θ even though β is non-zero. We can not determine θ dynamically in this case. At this number of flavour, $W_A = (A_{\hat{\mu}}, \lambda)$ and $Q^{adj} = (\phi_q^{adj}, q^{adj})$ form a massless vector multiplet of $N = 2$ super Yang-Mills theory in the 4-dimensions. Note that the effective potential corresponds to massless limit of the matters. In $N = 2$ supersymmetry there exists global $SU(2)_R$ symmetry in addition to the usual $U(1)_R$ symmetry. Under $SU(2)_R$ transformation, λ is interchanged into q^{adj} and vice versa. The λ and ϕ_q^{adj} can form $N = 1$ chiral superfield $Q^{adj} = (\lambda, \phi_q^{adj})$ instead of the original $N = 1$ chiral superfield $Q^{adj} = (q^{adj}, \phi_q^{adj})$ by using $SU(2)_R$ symmetry. The λ and ϕ_q^{adj} satisfy the same boundary condition as seen from (30) and (35). Hence, $N = 1$ supersymmetry remains to be unbroken though $N = 2$ supersymmetry is broken by the boundary conditions. The non-renormalization theorem works by this unbroken $N = 1$ supersymmetry, so that the effective potential vanishes even for non-zero values of β .

Finally, let us study the gauge symmetry breaking if we add both matters in the fundamental representation and those in the adjoint representation. The total effective potential is given by

$$\begin{aligned}\bar{V}_{eff}^{total} &= \bar{V}_{eff}^{SQCD} + \bar{V}_{eff}^{fd} + \bar{V}_{eff}^{adj} \\ &= -2F(t) + G(t, \beta) + N_F \left[-2G(t/2, \beta) + 4F(t/2) \right] \\ &\quad + N_F^{adj} \left[-G(t, \beta) + 2F(t) \right] \\ &= \frac{\beta^2}{12} \left[-3t^3 + 6\pi t - 2\pi^2 - \frac{1}{2}\beta^2 + N_F \left(\frac{3}{2}t^2 - 6\pi t + 4\pi^2 + \beta^2 \right) \right. \\ &\quad \left. + N_F^{adj} \left(3t^2 - 6\pi t + 2\pi^2 + \frac{1}{2}\beta^2 \right) \right].\end{aligned}$$

For $(N_F, N_F^{adj}) = (N_F, 0)$, $(N_F, 1)$, $SU(2)$ gauge symmetry is not broken because U_g^{inv} commutes with all the generators of $SU(2)$ in these cases. Note that we can not determine θ for $(N_F, N_F^{adj}) = (0, 1)$ because of the vanishing effective potential due to the unbroken $N = 1$ supersymmetry. Except for these cases the minimum of the total potential is located at

$$\theta = \frac{N_F + N_F^{adj} - 1}{N_F + 2N_F^{adj} - 2} \pi, \quad \text{for } 0 \leq \theta \leq \pi.$$

We obtain

$$U_g^{inv} = \cos\left(\frac{N_F + N_F^{adj} - 1}{N_F + 2N_F^{adj} - 2}\pi\right) + i\sigma^3 \sin\left(\frac{N_F + N_F^{adj} - 1}{N_F + 2N_F^{adj} - 2}\pi\right), \quad (37)$$

which commutes with only the third component of the $SU(2)$ generators T^3 . Therefore, $SU(2)$ gauge symmetry is broken to $U(1)$.

Let us study asymptotic behavior of the non-integrable phase θ . For fixed N_F , $\theta \rightarrow \pi/2$ as $N_F^{adj} \rightarrow \infty$, so that $U_g^{inv} \rightarrow -i\sigma^3$. The $SU(2)$ gauge symmetry is broken to $U(1)$ in this limit. While, for fixed N_F^{adj} , $\theta \rightarrow \pi$ as $N_F \rightarrow \infty$, so that $U_g^{inv} = -\mathbf{1}_{2 \times 2}$. The $SU(2)$ gauge symmetry is restored in this limit. When $N_F = N_F^{adj} \rightarrow \infty$, θ approaches to $2\pi/3$. The U_g^{inv} takes the values of $-\cos \pi/3 - i\sigma^3 \sin \pi/3$ in the limit. Hence, $SU(2)$ gauge symmetry is broken to $U(1)$ in this limit. We depict how the gauge symmetry breaking depends on number and the representation under the gauge group of the matters in Figure 1.

3.1.4 Effective Lagrangian on M^3

Taking into account of the discussions we have made, we obtain the effective lagrangian in the 3-dimensions. We begin from the $N = 1$ SQCD in the 4-dimensions and compactify it on $M^3 \otimes S^1$ with the fields having the boundary conditions (30) and (8) (or (35)). Then, the effective lagrangian on M^3 takes the form of

$$\mathcal{L}_{SQCD+matters}^{N=1} \longrightarrow \mathcal{L}_{SQCD+matters}^{eff} = \tilde{\mathcal{L}}_{SQCD+matters}^{N=2} + \tilde{\mathcal{L}}^{soft}. \quad (38)$$

The $\tilde{\mathcal{L}}_{SQCD+matters}^{N=2}$ is the lagrangian of $N = 2$ supersymmetric gauge theory with $SU(2)$ or $U(1)$ gauge group in the 3-dimensions. In the 3, 4-dimensional Minkowski space-time, the dimensions of the Dirac spinor are $2^{[D/2]} = 2, 4$, respectively. From the point of view of the 3-dimensions, $N = 1$ supersymmetry in the original 4-dimensions raises to $N = 2$ supersymmetry in the 3-dimensions by the dimensional reduction. The $N = 2$ supersymmetry in our effective theory consists of one vector multiplet and one (two) hyper multiplet(s), which are resulted from $N = 1$ spinorial chiral superfield W_A and $N = 1$ chiral superfield $Q^{adj}(Q, \overline{Q})$, respectively by the dimensional reduction. The $SU(2)$ gauge symmetry can be broken to $U(1)$ through the dynamics of the non-integrable phases, depending on number and the representation under the gauge group of the matters.

The $\tilde{\mathcal{L}}^{soft}$ stands for the soft supersymmetry breaking terms due to the boundary condition associated with the $U(1)_R$ symmetry. Its explicit form is obtained in (10). If the matters belong to the adjoint representation under $SU(2)$ gauge group, $(T^a)_{bc} = if_{bac}$, while if they belong to the fundamental representation, $T^a = \sigma^a/2$, where σ^a is the Pauli matrices. It should be understood that if the gauge symmetry is broken to $U(1)$, that is, $A_3^3 = \Phi^3$ takes the non-zero vacuum expectation values, then, physical field of Φ^a is the fluctuation around the vacuum expectation value. The effective lagrangian $\mathcal{L}_{SQCD+matters}^{eff}$ is the softly broken $N = 2$ supersymmetric gauge theory.

4 Conclusions and Discussions

We have discussed the effects of the boundary conditions of the fields for the compactified directions on the supersymmetric gauge theory. The supersymmetry can be broken explicitly by the boundary condition associated with the $U(1)_R$ symmetry. The effect of the breaking become manifest by redefining the fields so as to satisfy the periodic boundary condition. They turn out to appear as the soft supersymmetry breaking terms in the lagrangian. The effects are always soft supersymmetry breaking because they are generated only through the derivative ∂_μ in the kinetic terms, where the gauged $U(1)_R$ transformation by $U_R(y)$ is not respected as the symmetry of the theory. Remarkable feature is that the soft supersymmetry breaking terms do not have many arbitrary parameters, but they depend on an unique parameter β and the gauge coupling. The soft supersymmetry breaking terms are common to all matters, which are needed to avoid the FCNC. The supertrace of the squared mass matrix does not vanish because the supersymmetry is broken explicitly in our case. It should be stressed that these desirable soft supersymmetry breaking terms are automatically incorporated into the theory by the boundary condition associated with the $U(1)_R$ symmetry.

We have also discussed the effects of the boundary condition associated with the global gauge symmetry. The effects are interpreted as the constant background gauge field, which are dynamical degrees of freedom called non-integrable phases for the gauge field along the compactified direction in a multiply-connected space. Unlike the famous Aharanov-Bohm effects[15], the non-integrable phases are determined dynamically. We have shown explicitly that the gauge symmetry can be broken through the dynamics of the non-integrable phases in the SQCD. The gauge symmetry breaking depends on numbers and the representation under the gauge group of the matters.

We have begun from the $N = 1$ SQCD in the 4-dimensions and compactify it on $M^3 \otimes S^1$ with the fields having the boundary conditions for the S^1 -direction. As the result of the effects of the boundary conditions, we have finally obtained the effective lagrangian on M^3 , which is the softly broken $N = 2$ supersymmetric gauge theory. We can discuss both gauge and supersymmetry breaking in one scheme, say, the boundary conditions of the fields for the compactified directions.

We have evaluated the effective potential for the non-integrable phases (34), (36) in the massless limit of the matters. The gauge symmetry breaking through the dynamics of non-integrable phases are essentially caused by the infrared dynamics of the theory. The compactness of the extra coordinate S^1 shifts the zero point energies for massless particles, so that the gauge symmetry breaking is induced through the Casimir effect. If particles are massive, the gauge symmetry breaking may be different from the result obtained here[17]. It is interesting to study how the massive particle affect the gauge symmetry breaking in the supersymmetric gauge theory. We are studying the effects

of the boundary conditions on more realistic higher dimensional supersymmetric gauge theories. We believe that there are new possibilities for exploring models of softly broken supersymmetric gauge theories.

Acknowledgment

The author would like to thank Professor C. S. Lim for fruitful discussions, encouraging me. This work was supported by Grant-in-Aid for Scientific Research Fellowship, No.5106.

Appendix

In this appendix we present explicit expressions which are omitted in the subsection **2.3**. The 4-dimensional lagrangian for the massive matters is given by

$$\mathcal{L}_{matters} = \sum_I \left[Q_I^\dagger e^{2gV(N)} Q_I + \bar{Q}^{I\dagger} e^{2gV(\bar{N})} \bar{Q}^I \right]_D + W(Q, \bar{Q})_F,$$

where the subscripts F, D mean F, D -term, respectively, and $W(Q, \bar{Q}) \equiv m_I \bar{Q}^I Q_I$. The g is the gauge coupling constant, and $I (= 1, \dots, N_F)$ stands for flavour index. The $V(N) \equiv V^a T^a, (V(\bar{N}) \equiv V^a \bar{T}^a)$ is the vector superfield in the Wess-Zumino gauge. The $T^a (a = 1, \dots, N^2 - 1)$ is the generator of the gauge group, and $\bar{T}^a \equiv -T^{aT} = -T^{a*}$. Under the supersymmetric transformations δ_ξ defined by

$$\begin{aligned} \delta_\xi \phi_{qI} &= \sqrt{2} \xi q_I, & \delta_\xi \phi_{\bar{q}}^I &= \sqrt{2} \xi \bar{q}^I, \\ \delta_\xi q_I &= \sqrt{2} \xi F_{qI} + \sqrt{2} i \sigma^{\hat{\mu}} \bar{\xi} D_{\hat{\mu}} \phi_{qI}, & \delta_\xi \bar{q}^I &= \sqrt{2} \xi F_{\bar{q}}^I + \sqrt{2} i \sigma^{\hat{\mu}} \bar{\xi} D_{\hat{\mu}} \phi_{\bar{q}}^I, \\ \delta_\xi F_{qI} &= \sqrt{2} i \bar{\xi} \bar{\sigma}^{\hat{\mu}} D_{\hat{\mu}} \phi_{qI} + 2g \bar{\xi} \bar{\lambda} \phi_{qI}, & \delta_\xi F_{\bar{q}}^I &= \sqrt{2} i \bar{\xi} \bar{\sigma}^{\hat{\mu}} D_{\hat{\mu}} \phi_{\bar{q}}^I + 2g \bar{\xi} \bar{\lambda} \phi_{\bar{q}}^I, \end{aligned}$$

the lagrangian varies as $\mathcal{L}_{matters} = \partial_{\hat{\mu}} X_{matters}^{\hat{\mu}}$, where $X_{matters}^{\hat{\mu}} \equiv X_Q^{\hat{\mu}} + X_{\bar{Q}}^{\hat{\mu}}$ is calculated as

$$\begin{aligned} X_Q^{\hat{\mu}} &= \sum_I \left[ig \xi \sigma^{\hat{\mu}} \bar{\lambda}^a (\phi_{qI}^\dagger T^a \phi_{qI}) + \frac{i}{\sqrt{2}} \xi \sigma^{\hat{\mu}} \bar{q}^I F_{qI} + \frac{1}{\sqrt{2}} \xi \sigma^{\hat{\mu}} \bar{\sigma}^{\hat{\nu}} q_I (D_{\hat{\nu}} \phi_{qI})^\dagger \right. \\ &\quad \left. + \sqrt{2} i m_I \bar{\xi} \bar{\sigma}^{\hat{\mu}} q_I \phi_{\bar{q}}^I \right] + h.c. \end{aligned}$$

and

$$\begin{aligned} X_{\bar{Q}}^{\hat{\mu}} &= \sum_I \left[ig \xi \sigma^{\hat{\mu}} \bar{\lambda}^a (\phi_{\bar{q}}^{I\dagger} \bar{T}^a \phi_{\bar{q}}^I) + \frac{i}{\sqrt{2}} \xi \sigma^{\hat{\mu}} \bar{q}^I F_{\bar{q}}^I + \frac{1}{\sqrt{2}} \xi \sigma^{\hat{\mu}} \bar{\sigma}^{\hat{\nu}} \bar{q}^I (D_{\hat{\nu}} \phi_{\bar{q}}^I)^\dagger \right. \\ &\quad \left. + \sqrt{2} i m_I \bar{\xi} \bar{\sigma}^{\hat{\mu}} \bar{q}_I \phi_{qI} \right] + h.c. \quad . \end{aligned}$$

The lagrangian $\mathcal{L}_{matters}$ contains the kinetic terms for the quarks and the squarks. Their covariant derivatives are given by

$$\begin{aligned} D_{\hat{\mu}}\phi_{qI} &= \partial_{\hat{\mu}}\phi_{qI} - igT^a A_{\hat{\mu}}^a \phi_{qI}, & D_{\hat{\mu}}q_I &= \partial_{\hat{\mu}}q_I - igT^a A_{\hat{\mu}}^a q_I, \\ D_{\hat{\mu}}\phi_q^I &= \partial_{\hat{\mu}}\phi_q^I + ig\phi_q^I T^a A_{\hat{\mu}}^a, & D_{\hat{\mu}}\bar{q}^I &= \partial_{\hat{\mu}}\bar{q}^I + ig\bar{q}^I T^a A_{\hat{\mu}}^a. \end{aligned}$$

The derivative $\partial_{\hat{\mu}}$ in the covariant derivatives do not respect the symmetry of the theory under the gauged $U(1)_R$ transformation by $U_R(y) = e^{i\frac{\beta}{L}y}$. The supersymmetry breaking terms are generated only through the derivative as discussed in the text.

References

- [1] M. B. Green, J. H. Schwartz and E. Witten, *Superstring Theory*, Cambridge University Press 1987.
- [2] For a review, see for example, D. Bailin and A. Love, Rep. Prog. Phys. **50** (1987) 1087.
- [3] Y. Hosotani, Phys. Lett. **B126** (1983) 309.
- [4] Y. Hosotani, Ann. Phys. **190** (1989) 233.
- [5] K. Takenaga, hep-th/9710058.
- [6] See for example, J. Wess and J. Bagger, *Supersymmetry and Supergravity*, Princeton University Press 1992.
- [7] J. Scherk and J.H. Schwartz, Phys. Lett. **B82** (1979) 60.
- [8] C. Bachas, hep-th/9503030, references there in.
- [9] In this evaluation we assume that the gauge symmetry is not broken. Even if the gauge symmetry is broken, $\text{Str}\mathcal{M}^2$ does not vanish in our model.
- [10] D. J. Toms, Phys. Lett. **B126** (1983) 445.
- [11] A. Higuchi and L. Parker, Phys. Rev. **D37** (1988) 2853.
- [12] J. E. Hetrick and C. L. Ho, Phys. Rev. **D40** (1989) 4085.
- [13] This form of the constant background gauge field follows from assuming the diagonal form of U_g . (See (19)). But we restrict ourself to flat connection $\langle F_{\hat{\mu}\hat{\nu}} \rangle = 0$, so that the form of $\langle A_a \rangle$ on the torus is highly constrained such that all the components must commute. This yields the diagonal form of $\langle A_a \rangle$. On the other hand, if we assume the compactification on $M^{D-1} \otimes S^1$, we can always diagonalize U_g as $U_g =$

$C \text{diag}(e^{i\lambda_1}, \dots, e^{i\lambda_N}) C^\dagger$, where $C \in SU(N)$. In this case we can also write $\langle A_a \rangle$ as (26).

- [14] This expression corresponds to the effective potential for massless (s)quarks. In this calculation we have taken the massless limit of the matters. The $U(1)_R$ symmetry is not exactly defined in the limit, but the similar non-trivial phases are allowed from the single-valueness of the lagrangian even if we begin from the massless (s)quarks.
- [15] Y. Aharanov and D. Bohm, Phys. Rev. **D115** (1959) 484.
- [16] L. Brink, J. H. Schwartz and J. Scherk, Nucl. Phys. **B121** (1977) 77.
- [17] Y. Hosotani, Phys. Lett. **B129** (1983) 193.




Fig. 1

This figure shows how the gauge symmetry breaking depends on number and the representation under $SU(2)$ gauge group of the matters. The $\otimes$ stands for the gauge symmetry breaking to $U(1)$. On the other hand, $\circ$ stands for the unbroken $SU(2)$ symmetry. At the point $(N_F, N_F^{adj}) = (0, 1)$ denoted by $\times$ in the figure, we have unbroken $N = 1$ supersymmetry. Details are discussed in the text.